

OFFICE OF NAVAL RESEARCH


GRANT or CONTRACT: N00014-93-1-0772

R & T Code 313T006

Technical Report #10

Tetramethylammonium Phenyltrialkylborates in the Photoinduced Electron Transfer Reaction with Benzophenone. Generation of Alkyl Radicals and Their Addition to Activated Alkenes.

Polykarpov, A.; ; Neckers, D. C. *Tetrahedron Letters*, 1995 in press.



D. C. Neckers
Center for Photochemical Sciences
Dept. of Chemistry
Bowling Green State University
Bowling Green, OHIO 43402

Date Submitted
June 15, 1995

Reproduction in whole, or in part, is permitted for any purpose of the United States Government.

This document has been approved for public release and sale, its distribution is unlimited.

19950705 069

DTIC QUALITY INSPECTED 5

REPORT DOCUMENTATION PAGE


OMB No. 0704-0188

1. The reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204 Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188) Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE 6-15-95		3. REPORT TYPE AND DATES COVERED Technical Report	
4. TITLE AND SUBTITLE Tetramethylammonium Phenyltrialkylborates in the Photoinduced Electron Transfer Reaction with Benzophenone. Generation of Alkyl Radicals and Their Addition to Activated Alkenes.				5. FUNDING NUMBERS G N00014-93-J-1921 R & T 313T006	
6. AUTHOR(S) D.C. Neckers				Kenneth Wynne	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Center for Photochemical Sciences Bowling Green State University Bowling Green, Ohio 43402				8. PERFORMING ORGANIZATION REPORT NUMBER 10	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Department of the Navy Office of Naval Research 800North Quincy Street Arlington, VA 22217-5000				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Polykarpov, A.; Neckers, D. C. <i>Tetrahedron Letters</i> , 1995 in press.					
12a. DISTRIBUTION/AVAILABILITY STATEMENT Reproduction in whole or in part is permitted for any purpose of the United States Government. This document has been approved for public release and sale; its distribution is unlimited.				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Photoinduced one electron oxidation of tetramethylammonium phenyltrialkylborates by the excited state of benzophenone in an acetonitrile/benzene solution containing an excess of activated alkene produces substantially more than one equivalent of the alkyl radicals. The corresponding adducts of alkyl radicals to the alkenes are produced in good yields. No phenyl radical adducts are observed.					
14. SUBJECT TERMS				15. NUMBER OF PAGES	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified		18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified		19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	
				20. LIMITATION OF ABSTRACT UL	

Tetramethylammonium Phenyltrialkylborates in the Photoinduced Electron Transfer Reaction with Benzophenone. Generation of Alkyl Radicals and Their Addition to Activated Alkenes.

Alexander Y. Polykarpov and Douglas C. Neckers*
Center for Photochemical Sciences, Bowling Green State University, Bowling Green, OH 43403, USA.



Photoinduced one electron oxidation of tetramethylammonium phenyltrialkylborates by benzophenone triplet leads to formation of substantially more than one equivalent of alkyl radicals. The adducts of alkyl radicals to various electron poor alkenes are produced in good yields. No phenyl radical adducts are observed.

Accession For	NTIS CRA&I	<input checked="" type="checkbox"/>	DTIC TAB	<input type="checkbox"/>	Unannounced	<input type="checkbox"/>	Justification	By	Distribution /	Availability Codes	Dist	A-1
											Avail and/or Special	

Tetramethylammonium Phenyltrialkylborates in the Photoinduced Electron Transfer Reaction with Benzophenone. Generation of Alkyl Radicals and Their Addition to Activated Alkenes.

Alexander Y. Polykarpov and Douglas C. Neckers*

Center for Photochemical Sciences¹


Bowling Green State University

Bowling Green, Ohio 43403.

Abstract: Photoinduced one electron oxidation of tetramethylammonium phenyltrialkylborates by the excited state of benzophenone in an acetonitrile/benzene solution containing an excess of activated alkene produces substantially more than one equivalent of the alkyl radicals. The corresponding adducts of alkyl radicals to the alkenes are produced in good yields. No phenyl radical adducts are observed.

Trialkylboranes are known to produce alkyl group addition products to certain olefins under various conditions.² Though radical mechanisms are suggested, in the majority only one alkyl group of three may be used in subsequent reaction processes. Unlike α,β -unsaturated ketones and aldehydes, α,β -unsaturated esters and nitriles do not readily undergo addition reactions with organoboranes by most of the reported processes³. Though it has been reported that ethyl esters of acrylic and methacrylic acids react with organoboranes to form conjugate addition products under electrolytic conditions, less than one equivalent of alkyl group was transferred from the boranes.⁴

Photochemical oxidation of triarylalkylborates has been subject of intensive activity in the open and patent literature since cyanine borates were first discovered to initiate acrylate photopolymerization.⁵ Trialkylaryl borates, on the other hand, have been much less extensively studied. In this communication we describe preliminary studies of the addition of primary and secondary alkyl radicals generated by single electron transfer from tetramethylammonium phenyltrialkylborates to benzophenone triplet to various activated alkenes (eq. 1). In contrast to the other systems,² substantially more than one equivalent of alkyl radicals may be generated using our reaction, and the reaction is general. Efficient radical additions are reported for a variety of electron poor olefins.



The addition of phenyl anion to trialkylboranes, available from alkenes (method A) or reaction of dichlorophenylboron with three equivalents of the alkyl anion (method B) followed by cation exchange (in water for most borates except those containing C₆ and longer hydrocarbon chains) gives various tetramethylammonium phenyltrialkylborates in good yields (eq. 2). Most borate salts are water insoluble

solids, relatively stable to air and moisture. Acetonitrile solutions of these borates show no absorbance above 300nm. Yields of tetramethylammonium phenyltrialkylborates obtained by these methods and ^{11}B nmr data are given in Table 1. The X-ray diffraction structure shows tetramethylammonium phenyltriethylborate to be tetrahedral (Figure 1).





Table 1. Yields^a and ^{11}B NMR Data for Tetramethylammonium Phenyltrialkylborates ($\text{PhBR}_3\text{NMe}_4$).

Entry	R	Yield (%)	δ (ppm)	Entry	R	Yield (%)	δ (ppm)
1	Methyl ^b	70	-33.89 ^c	6	Cyclopentyl	90	-32.44
2	Ethyl	95	-33.07	7	Cyclohexyl	75	-31.60
3	sec-Butyl	82	-30.16	8	Cycloheptyl	76	-29.36
4	n-Hexyl	68	-33.96	9	Norbornyl	95	-31.35
5	n-Octyl	61	-34.04	10	n-Bu ₄ BNMe ₄ ^d	92	-36.86

^aOverall yields. ^bObtained through method B from methyllithium (other borates were made via procedure A). ^cIn d₆-DMSO, $\delta(\text{B}(\text{OMe})_3)=0$ ppm. ^dTetramethylammonium tetra-n-butylborate, readily available from tri-n-butylborane and n-butyllithium, is given for comparison. It possesses good reactivity in the radical addition. As with other tetraalkylborates, it is less stable than the phenyltrialkylborates.

Figure 1. Molecular Structure of Tetramethylammonium Phenyltriethylborate (only the anion is shown).



Irradiation (Rayonet, 350nm, water cooling, 25°C, 2-4 hours) of a solution (under nitrogen) containing borate (5mM), benzophenone (2-5mM) and an activated alkene (50mM) in benzene/acetonitrile (5:1 v/v) gives adducts in good yields. Yields of the adducts of ethyl radicals generated from tetramethylammonium phenyltriethylborate to a variety of activated alkenes are presented in Table 2.

Table 2. Yields of Adducts^a of Ethyl Radicals Generated From Tetramethylammonium Phenyltriethylborate to Various Activated Alkenes.

Entry	Alkene	Adduct	Yield (%)
1	n-Butyl acrylate ^b	n-Butyl pentanoate	190 ^c
2	Methyl methacrylate ^b	Methyl 2-methylpentanoate	184
3	Phenyl vinylsulfone	Phenyl n-butylsulfone	(168) ^d
4	Diethyl maleate	Diethyl ethylsuccinate	(160)
5	Diethyl vinylphosphonate	Diethyl n-butylphosphonate	(153)
6	Acrylonitrile ^b	Valeronitrile	(145)
7	N,N-Dimethyl acrylamide	N,N-Dimethyl valeroamide	119
8	Styrene	n-Butylbenzene	73

^aAll the products were either compared with authentic samples or exhibited analytical data and spectra in accordance with the assigned structures. Yields are based on borate used. ^bUsed with inhibitor. ^cDetermined by GC. ^dIsolated Yields.

The results of the addition of alkyl radicals generated from tetramethylammonium phenyltrialkylborates to n-butyl methacrylate are given in Table 3. Though the mechanism of the reaction is not completely understood, and lithium phenyltrialkylborates have been found to be significantly less reactive (sometimes completely unreactive) than the corresponding tetramethylammonium salts under the conditions discussed above, the initial step is anticipated to be an electron transfer from borate to benzophenone triplet state followed by formation of the alkyl radical and phenyldialkylborane.⁵ Alkyl radicals form because they are of greater stability and phenyl radical products (biphenyl, phenyl adducts) were not detected by GC (less than 0.1%). Such alkyl radicals, when formed, add to the alkene. Phenyldialkylborane is also capable of initiating a chain radical reaction. It can react either with the oxygen centered radical product of addition of the alkyl radical to the alkene or with the triplet state of benzophenone^{3,6} forming additional radicals. Benzophenone is usually recovered (10-40%). Water soluble and polar boron containing products have yet to be studied.

The preliminary results outlined in this communication indicate that photoinitiated oxidation of tetramethylammonium phenyltrialkylborates, easily accessible from alkenes via hydroboration followed by the addition of phenyl anion and further cation exchange, leads to formation of substantially more

than one equivalent of the adducts. Further studies are under way to define the mechanism more clearly and to extend these investigations to other systems involving different excited state acceptors, alkenes and various borates.

Table 3. Yields of Adducts^a in the Addition of Primary and Secondary Alkyl Radicals Generated From Various Tetramethylammonium Phenyltrialkylborates to n-Butyl Methacrylate^b.

Entry	R	Adduct	Yield (%)
1	Methyl	n-Butyl 2-methylbutanoate	90
2	Ethyl	n-Butyl 2-methylpentanoate	193
3	sec-Butyl	n-Butyl 2,4-dimethylhexanoate	236
4	n-Octyl	n-Butyl 2-methylundecanoate	(221) ^c
5	n-Hexyl	n-Butyl 2-methylnonanoate	(169)
6	Cyclopentyl	n-Butyl 3-cyclopentyl-2-methylpropanoate	(240)
7	Cyclohexyl	n-Butyl 3-cyclohexyl-2-methylpropanoate	(175)
8	Cycloheptyl	n-Butyl 3-cycloheptyl-2-methylpropanoate	(185)
9	Norbornyl	n-Butyl 3-norbornyl-2-methylpropanoate	(227)

^aYields are based on borate consumed and determined by GC. ^bUsed with inhibitor. ^cYields were determined after isolation by flash chromatography (silica gel, hexanes / ethyl acetate 40/1).

Acknowledgments: This work is supported, in part, by a Fellowship from the McMaster Endowment (A.Y.P.), by the N.S.F. (DMR 9013109), and by O.N.R. (N00014-93-1-0772). We also wish to thank Dr. A. A. Pinkerton and A. Martin from the University of Toledo for their assistance with x-ray diffraction analysis.

References.

1. Contribution #222 from the Center for Photochemical Sciences.
2. Brown, H.C. *Organic Synthesis via Boranes*; John Wiley & Sons, Inc., New York, **1975**, pp 146-149 and references cited therein.
3. Wilkinson, G.; Stone, F.G.A.; Abel, E.W. *Comprehensive Organometallic Chemistry. The Synthesis, Reactions and Structures of Organometallic Compounds*; Pergamon Press, Oxford, **1982**, vol. 7, pp. 291-293, 335 and references cited therein.
4. Takahashi, Y.; Yuasa, K.; Tokuda, M.; Itoh, M.; Suzuki, A. *Bull. Chem. Soc. Japan* **1978**, 51, 339-340.
5. See for example: Schuster G.B. *Pure & Appl. Chem.* **1990**, 62, 1565-1572. Shanklin, M.S.; Gottschalk, P.; Adair, P.C. US Patent 5055372 (Prior. 10. 8. 91) to Mead Corporation.
6. Davies, A.G.; Scaiano, J.C. *J. Chem. Soc. Perkin Trans. 2* **1972**, 2234-2238.